

On the relation between the boson-fermion model and RVB state

Evgueny Kochetov

Bogolyubov Theoretical laboratory, Joint Institute for Nuclear Research, 141980 Dubna, Russia

Marcin Mierzejewski

Institute of Physics, University of Silesia, 40-007 Katowice, Poland

We show that the resonating valence bond (RVB) state of the t - J Hamiltonian can be described within an effective boson-fermion model. We derive a boson-fermion Hamiltonian which describes the lowest order coupling between holons and spinon waves. The RVB order parameter determines a hybridization between the both types of excitations. The effective Hamiltonian is investigated within the lowest order self-consistent conserving diagrammatic approximation. We compare our approach with the phenomenological boson-fermion model (BFM) that is believed to describe the pseudogap phase.

74.20.Mn, 75.10.Jm

The origin of the pseudogap is one of the most important current problems in high temperature superconductors. This normal state gap is seen by various techniques, including angle-resolved photoemission spectroscopy [1–3], NMR [4,5], infrared [6], and transport [7] measurements. Although we now possess considerable experimental data, the understanding of this phenomenon is still far from complete. There are a number of microscopic scenarios, but none of them is commonly accepted. The opening of the pseudogap, and its evolution into the true superconducting gap, may, however, be described by a phenomenological boson-fermion model [8–12]. This BFM is usually understood as a system of localized tightly bound electron pairs (bosons) which hybridize with itinerant electrons. This model customarily neglects the hard core effects of the bound electron pairs [9–11] and assumes the standard commutation relations for the boson operators. We may then identify three characteristic temperatures [10] which correspond to: (i) the opening of the pseudogap in the fermion excitation spectrum, (ii) the appearance of itinerant bosons, (iii) the condensation of these bosons and the concurrent onset of superconductivity. Due to the phenomenological character of the BFM, the microscopic origin of the pair correlations is not specified. Moreover, it is not straightforward to select an appropriate set of model parameters — in particular the total number of fermions and bosons. In order to answer these questions we need to find a relationship between the BFM and a microscopic approach [13]. The aim of this work is to provide such a link by demonstrating a close similarity between the BFM and the t - J model with a mean-field RVB state.

As it is known, the RVB state [14] is characterized by three important features: (i) spin-charge separation,

(ii) short range antiferromagnetic (AF) correlations, and (iii) holon condensation. Owing to (i), the RVB pairing amplitude may develop at a much higher temperature than the superconducting transition temperature, at which the phase coherence between the electron pairs is established [15]. In the RVB scenario [14] there is a phase with the preformed, incoherent, electron pairs with an amplitude Δ_{RVB} that serves as the RVB order parameter. In this phase the RVB electron pairs are to be thought of as spinon singlets which are coupled to holon excitations, this resulting in a boson-fermion (BF) interaction. It is therefore natural to suggest that it is the RVB order parameter that controls the pseudogap. This is our basic assumption.

First we demonstrate that the RVB state can be described by a holon-spinon interaction. Next, application of the linear spinon-wave approximation leads to a boson-fermion model where bosons are doubly charged with respect to spinless fermions.

To begin, we express the t - J model in terms of the Hubbard operators $X_i^{\sigma 0} = c_{i\sigma}^\dagger(1 - n_{i,-\sigma})$,

$$H_{t-J} = -t \sum_{\langle i,j,\sigma \rangle} X_i^{\sigma 0} X_j^{0\sigma} + \text{H.c.} + J \sum_{\langle i,j \rangle} \left(\vec{Q}_i \vec{Q}_j - \frac{n_i n_j}{4} \right),$$

where $c_{i\sigma}$ annihilates an electron at site i with spin projection $\sigma = \pm$, $n_{i\sigma} \equiv c_{i\sigma}^\dagger c_{i\sigma}$, $n_i = \sum_{\sigma} n_{i\sigma}$ and $\sum_{\langle i,j \rangle}$ denotes a summation over the nearest neighbor *nonrepeated* bonds. The electron-spin operator, $\vec{Q}_i = \frac{1}{2} \sum_{\sigma\sigma'} X_i^{\sigma 0} \vec{\tau}_{\sigma\sigma'} X_i^{0\sigma'}$, enters the magnetic part of H_{t-J} and $\vec{\tau} = (\tau^1, \tau^2, \tau^3)$ stand for the Pauli matrices. The operators $\vec{Q}$ fulfil the SU(2) algebra commutation relations, but do not form a true representation, since the second Casimir operator $\vec{Q}^2$ is not a c-number.

Note that the $X^{\sigma 0}$ appear as fermionic-like operators, while the $X^{\sigma\sigma'} = X^{\sigma 0} X^{0\sigma'}$ correspond to bosonic degrees of freedom. The crucial t - J model requirement of no double occupancy $\sum_{\sigma} c_{i\sigma}^\dagger c_{i\sigma} \leq 1$ follows automatically: by definition $X^{00} + \sum_{\sigma} X^{\sigma\sigma} = I$. Unlike the slave-particle representations, this restriction imposes an additional constraint on dynamical degrees of freedom and seems to pose a severe problem in a mean-field treatment. Note further that the t - J Hamiltonian possesses two global U(1) symmetries: $U_{N_e}(1)$ and $U_{Q_3}(1)$. These correspond to the conservation of the total electron number $N_e = \sum_i (X_i^{++} + X_i^{--}) = \sum_i n_i$, and to the total spin projection $Q_3 = \frac{1}{2} \sum_i (X_i^{++} - X_i^{--})$, respectively.

A conventional Hartree-Fock decoupling is applied to the magnetic part of the t - J Hamiltonian

$$H_J := J \sum_{\langle i,j \rangle} \left(\bar{Q}_i \bar{Q}_j - \frac{1}{4} n_i n_j \right) = -J \sum_{\langle i,j \rangle} b_{ij}^\dagger b_{ij}$$

to yield

$$H_J \simeq -J \sum_{\langle i,j \rangle} (\Delta_{ij} b_{ij}^\dagger + \Delta_{ij}^* b_{ij} - |\Delta_{ij}|^2),$$

where $b_{ij}^\dagger = \frac{1}{\sqrt{2}} (X_i^{+0} X_j^{-0} - X_i^{-0} X_j^{+0})$, is the valence bond "singlet" pair creation operator and Δ_{ij} is the RVB order parameter defined on each bond between the nearest neighbor sites. $\Delta_{ij} \neq 0$ breaks the global $U_{N_e}(1)$ symmetry, though it does not directly result in superconductivity. It instead indicates the onset of the electron spin-singlet formation. There are many possible mean field states in RVB theory, among which the extended s -phase is selected here because of its simplicity.

To proceed, we employ the Feynman path-integral representation for the partition function. We use the *geometric quantization approach* based on the Poisson realization of the $SU(2|1)$ superalgebra on a classical phase space — the supersphere $CP^{1|1}$ [16]. In this approach there is an isomorphic map $X \rightarrow X^{cl}$ of the $SU(2|1)$ generators into classical functions on the supersphere which preserves the Lie superalgebra structure. This map can be realized explicitly by exhibiting covariant symbols (known as the momentum maps) of the Hubbard operators in terms of the $SU(2|1)$ coherent states, $|z, \xi\rangle$ [16]: $X^{cl} = \langle z, \xi | X | z, \xi \rangle$. Then, $Q_3^{cl} = -\frac{1}{2}(1 - |z|^2)w$, $(Q^+)^{cl} = zw$, $(Q^-)^{cl} = \bar{z}w$, $(X^{0-})^{cl} = -z\bar{\xi}w$, $(X^{0+})^{cl} = -\bar{\xi}w$, $(X^{+0})^{cl} = -\xi w$, $(X^{-0})^{cl} = -\bar{z}\xi w$, $(X^{++})^{cl} + (X^{--})^{cl} = (1 + \bar{z}z)w$, $(X^{00})^{cl} = \bar{\xi}\xi w$, where $w := (1 + |z|^2 + \bar{\xi}\xi)^{-1}$. Here the even z and odd ξ Grassmann variables parameterize superspace $SU(2|1)/U(1|1) = CP^{(1|1)}$, which is the $N = 1$ supersymmetric extension of the Riemannian sphere CP^1 . These functions form a $SU(2|1)$ superalgebra under the Poisson superbrackets on $CP^{1|1}$. For an arbitrary fermionic Hamiltonian with the requirement of no double occupancy, the relevant Berezin path-integral representation of the partition function is equivalent to the $SU(2|1)$ phase-space path integral

$$Z = \int_{CP^{1|1}} D\mu_{SU(2|1)} \exp \left(\int \langle z, \xi | \frac{d}{dt} - H | z, \xi \rangle dt \right),$$

$$D\mu_{SU(2|1)} = \frac{d\bar{z}dzd\bar{\xi}d\xi}{2\pi i(1 + |z|^2 + \bar{\xi}\xi)},$$

where the Hamiltonian is expressed in terms of the X -operators. Despite a rather complicated appearance, the $SU(2|1)$ path-integral representation for the t - J model is convenient since (i) it automatically incorporates the requirement of no double occupancy, $(X^{00})^{cl} +$

$\sum_\sigma (X^{\sigma\sigma})^{cl} \equiv I$, independently of any approximations being applied to z and ξ fields; (ii) it furnishes a convenient set of independent dynamical variables $(z, \xi) \in CP^{1|1}$, which can directly be related to the charge and spin degrees of freedom.

The coordinates z and ξ can be related to the $SU(2)$ spinon and $U(1)$ holon degrees of freedom in the following manner: We make two successive changes of variables $z \rightarrow z\sqrt{1 + \bar{\xi}\xi}$, $\xi \rightarrow \xi\sqrt{1 + |z|^2}$ to decouple the $SU(2|1)$ measure, $D\mu_{SU(2|1)} \rightarrow D\mu_{SU(2)} D\mu_{U(1)}$, where $D\mu_{SU(2)} = d\bar{z}dz/\pi i(1 + |z|^2)^2$ stands for the $SU(2)$ invariant measure on CP^1 , while $D\mu_{U(1)} = d\bar{\xi}d\xi$ denotes the Berezin integration over odd Grassmann variables [17].

The t - J partition function becomes

$$Z_{t-J} = \int_{CP^1} \prod_j D\mu_{SU(2)}^{(j)} \int \prod_j D\mu_{U(1)}^{(j)} \exp[\mathcal{A}_{t-J}],$$

$$\mathcal{A}_{t-J} = \frac{1}{2} \sum_j \int_0^\beta \left(\frac{\dot{\bar{z}}_j \dot{z}_j - \bar{z}_j \dot{\bar{z}}_j}{1 + |z_j|^2} + \dot{\xi}_j \xi_j - \bar{\xi}_j \dot{\xi}_j \right) dt - \int_0^\beta H_{t-J}^{cl} dt, \quad (1)$$

with ξ_i corresponding to the $U(1)$ charged spinless fermion degrees of freedom (holons), and z_i representing pure $SU(2)$ spins (spinons).

The classical Hamiltonian now reads

$$H_{t-J}^{cl} = -t \sum_{\langle i,j \rangle} \xi_i \bar{\xi}_j \langle z_i | z_j \rangle - \frac{J\Delta_{RVB}}{2} \sum_{\langle i,j \rangle} \xi_i \xi_j \Phi(\bar{z}_j, \bar{z}_i) + \text{H.c.} - \mu' \sum_i (1 - \bar{\xi}_i \xi_i) - \lambda \sum_i [2S_3(\bar{z}_i, z_i) + \bar{\xi}_i \xi_i]. \quad (2)$$

We have dropped the constant term. We have also explicitly introduced a chemical potential term, $-\mu' N_e$, as well as the Langrange multiplier λ which controls the magnitude of the total electron magnetic moment.

The $SU(2)$ algebra is realized on CP^1 in terms of the momentum maps $S_3^{cl} = -\frac{1}{2} \frac{1 - |z|^2}{1 + |z|^2}$, $(S^+)^{cl} = \frac{\bar{z}}{1 + |z|^2}$, $(S^-)^{cl} = \frac{z}{1 + |z|^2}$, $z \in CP^1$ and

$$\Phi(\bar{z}_j, \bar{z}_i) = \frac{(\bar{z}_j - \bar{z}_i)}{\sqrt{(1 + |z_i|^2)(1 + |z_j|^2)}} =: \Phi_{ij}$$

appears as a spinon-singlet amplitude, $\Phi_{ij} = \Psi_{s_3=-1/2}(z_i) \Psi_{s_3=1/2}(z_j) - \Psi_{s_3=-1/2}(z_j) \Psi_{s_3=1/2}(z_i)$, where $\Psi_{s_3=\pm 1/2}(z)$ is a spinon wave function in the basis of the $SU(2)$ coherent states, $|z\rangle = (1 + |z|^2)^{-1/2} \exp\{z S_-\}_{|s_3=1/2}$. Namely, $\Psi_{s_3=\pm 1/2}(z) = \langle s_3 = \pm 1/2 | z \rangle$, $S_{(3)}|s_3\rangle = s_3|s_3\rangle$ and the $SU(2)$ coherent-state inner product is

$$\langle z_i | z_j \rangle = \frac{(1 + \bar{z}_i z_j)}{\sqrt{(1 + |z_i|^2)(1 + |z_j|^2)}},$$

so that $|\Phi_{ij}|^2 = 1 - |\langle z_i | z_j \rangle|^2$.

In general, the Hamiltonian (2) cannot be thought of as a covariant symbol of a Fermi–Bose type operator, except for a particular case of the linear *spinon*–wave (LSW) approximation. This approximation amounts to expanding the action (1), as well as a measure factor in the path integral, in powers of $|z|^2$ up to the leading order. In the paramagnetic phase $\langle Q_{3,i}^{LSW} \rangle = 0$, and the spinon fluctuations become bounded by the condition $\langle |z_i|^2 \rangle \leq 1/2$. This, to some extent, justifies the LSW approximation. Within this approximation the partition function, takes on the form

$$Z_{\text{BF}} = \int \prod_j d\bar{z}_j dz_j d\bar{\xi}_j d\xi_j \exp[\mathcal{A}_{\text{BF}}],$$

$$\mathcal{A}_{\text{BF}} = \frac{1}{2} \sum_j \int_0^\beta \left(\dot{\bar{z}}_j z_j - \bar{z}_j \dot{z}_j + \dot{\bar{\xi}}_j \xi_j - \bar{\xi}_j \dot{\xi}_j - H_{\text{BF}}^{\text{cl}} \right) dt,$$

where the classical Hamiltonian reads:

$$H_{\text{BF}}^{\text{cl}} = -t \sum_{\langle i,j \rangle} \xi_i \bar{\xi}_j - \frac{J\Delta_{\text{RVB}}}{2} \sum_{\langle i,j \rangle} \xi_i \xi_j (\bar{z}_j - \bar{z}_i) + \text{H.c.}$$

$$- \mu' \sum_i (1 - \bar{\xi}_i \xi_i) - \lambda \sum_i [-1 + 2\bar{z}_i z_i + \bar{\xi}_i \xi_i].$$

It straightforward to see that the above partion function represents the following boson–fermion system:

$$H_{\text{BF}} = -t \sum_{\langle i,j \rangle} f_i f_j^\dagger - 2v \sum_{\langle i,j \rangle} f_i f_j (b_j^\dagger - b_i^\dagger) + \text{H.c.}$$

$$+ \Delta_B \sum_i b_i^\dagger b_i - \mu \sum_i (2b_i^\dagger b_i + f_i^\dagger f_i),$$

$$\{f_i, f_j^\dagger\} = [b_j, b_j^\dagger] = \delta_{ij}. \quad (3)$$

The parameters of the boson–fermion system are expressed in terms of the parameters of the original t – J Hamiltonian and the RVB state: $v = J\Delta_{\text{RVB}}/4$; $\Delta_B = 2\mu'$; $\mu = \mu' + \lambda$. The $U_{Q_3}(1)$ symmetry of eq. (2) under global gauge transformations $z_i \rightarrow e^{2i\theta} z_i$, $\xi_i \rightarrow e^{i\theta} \xi_i$ leads to conservation of the total charge $\sum_i (2b_i^\dagger b_i + f_i^\dagger f_i)$. Note, that this global $U(1)$ symmetry characterizes the standard BFM and indicates that bosons are doubly charged with respect to fermions. The BFM discussed in Refs. [9–11] is referred to as the standard approach. One should also keep in mind that this symmetry is an intimate feature of the RVB state, and occurs independently of the LSW approximation.

The standard BFM was investigated with the help of various methods, e.g.: the lowest order self-consistent conserving diagrammatic approximation [9,10], the dynamical mean-field approximation [12] and the continuous unitary transformation [11]. Unlike the standard BFM, the Hamiltonian obtained in the present study involves spinless fermions. Therefore decay and recombination between fermions and bosons involves two neighboring lattice sites. In the present case, bosons with momentum $\mathbf{q} = 0$ do not participate in these processes, in the

standard BFM these bosons are of crucial importance. Because of these differences, we cannot directly adopt the numerical results obtained for the standard BFM. Therefore, we have investigated a one dimensional (1D) case within the lowest order self-consistent conserving diagrammatic approximation and compared our results to those presented in Refs. [9,10].

In the the lowest order self-consistent conserving diagrammatic approximation one obtains a system of equations for the fermion, $G_F(\mathbf{k}, \omega)$, and boson, $G_B(\mathbf{k}, \omega)$, Green’s function. We refer to Ref. [10] for the details. This set of equations has been solved iteratively slightly above the real axis, i.e. for $\omega = \text{Re}\omega + i0.01t$. We have used grids of 501 momenta in the interval $[0, \pi]$, and 2001 frequencies in the interval $[-5t, 5t]$. We have also fixed the bosonic level $\Delta_B = -2.4t$.




FIG. 1. The fermionic density of states calculated at the temperature $kT = 0.02t$ for different values of the boson–fermion coupling. $\omega = 0$ corresponds the Fermi level. The inset shows the average bosonic occupation number as a function of momentum.

Fig. 1 shows the fermionic density of states, $\rho(\omega)$, calculated for different values of the RVB order parameter. The boson–fermion coupling enhances the density of states just above the Fermi level and reduces ρ for much larger energies. Effectively, the local minimum in the density of states occurs at the Fermi level. Modification of the density of states takes place also for a weak boson–fermion coupling, when the local minimum at the Fermi level is almost not visible. Therefore, despite the absence of the gap the system cannot be considered as a composition of noninteracting fermions and bosons. The most important difference between the standard BFM and the present study shows up in the average boson occupation number n_B (see the insets in Figs. 1 and 2). Here, the boson–fermion coupling enhances the number of bosons with momentum close to π , whereas in the standard BFM the bosons with momentum $\mathbf{q} = 0$ play the dominating role. In the present case the momentum dependence of n_B can be attributed to the AF short range correlations. However, bosons are uniformly distributed

in the real space and there is no antiferromagnetic order (in agreement with the assumption of the RVB state).




FIG. 2. The fermionic density of states calculated for (i) non interacting particles, (ii) standard BFM (fermions with spin) and (iii) obtained in the present study (spinless fermions). The inset shows the average boson occupation number as a function of momentum in the cases (ii) and (iii).

In Fig. 2 we compare $\rho(\omega)$ and n_B obtained within the standard BFM and the present study. In the case of spinless fermions $n_B(\mathbf{k})$ is relatively flat, when compared to the standard approach. It suggests that bosons do not condensate at the state with some particular momentum. However, investigation of this problem is not possible within a one-dimensional numerical study. One should also keep in mind, that in our approach (which concerns only the mean field RVB state) condensation of bosons does not represent the superconducting phase transition. Similarly to the standard BFM, the local minimum in ρ , that occurs at the Fermi level, deepens with the decreasing temperature and disappears when the temperature is sufficiently high (see Fig. 3).

The difference between the present approach and the standard BFM shows also up in the fermionic spectral functions (see Fig. 4). Here, the most important broadening of the spectral functions takes place above and below the Fermi surface. In the standard case [9] the spectral functions have a three-peaked structures, which are mostly visible at the Fermi surface. It may be attributed to the free-particle-like motion of the bosons with small wave vectors. However, in the present approach these bosons are not involved in the decay and recombination processes.




FIG. 3. The same as in Fig. 1 but calculated at different temperatures for $v = 0.15t$.

To conclude, starting from an RVB state for the t - J Hamiltonian, we have derived a connection between the t - J model and the BFM. We have expressed the parameters of the effective Hamiltonian in terms of those of the t - J model and the RVB state. The resulting model differs from the standard one in that the fermionic degrees of freedom correspond to spinless quasiparticles. However, as in the standard BFM, a local minimum in the fermionic density of states occurs close to the Fermi level. The minimum deepens with the decreasing of temperature and vanishes when the temperature is sufficiently high. The present study is restricted only to the mean-field RVB state, and does not attempt to account for the superconducting phase transition. It is, however, free from the ambiguities which occur in the phenomenological BFM when tries to select an appropriate set of model parameters.




FIG. 4. Spectral function for fermions calculated for wave-vectors, which are in the vicinity of the Fermi surface. We have used $v = 0.15t$ and $kT = 0.02t$.

- [1] D.S. Marshall *et al*, Phys. Rev. Lett., **76**, 4841 (1996).
- [2] H. Ding *et al*, Nature (London) **382**, 51 (1996).
- [3] M.R. Norman *et al*, Nature (London) **392**, 157 (1998).
- [4] G.V.M. Williams *et al*, Phys. Rev. Lett. **78**, 721 (1997).
- [5] G.V.M. Williams *et al*, Phys. Rev. Lett. **80**, 337 (1998).
- [6] D.N. Basov *et al*, Phys. Rev B**50**, 3511 (1994).
- [7] J.L. Tallon *et al*, Phys. Rev. Lett. **75** 4114 (1995).
- [8] S. Robaszkiewicz, R. Micnas, and J. Ranninger, Phys. Rev. B**36**, 180 (1987).
- [9] J. Ranninger, J. M. Robin, and M. Eschrig, Phys. Rev. Lett. **74**, 4027 (1995).
- [10] P. Devillard and J. Ranninger, Phys. Rev. Lett. **84**, 5200 (2000).
- [11] T. Domański and J. Ranninger, Phys. Rev. B**63**, 134505.
- [12] A. Romano, Phys. Rev. B**64**, 125101 (2001).
- [13] E. Altman and A. Auerbach, Phys. Rev. B**65**, 104508 (2002).
- [14] P.W. Anderson, Science **235**, 1196 (1987); G. Baskaran, Z. Zou and P.W. Anderson, Solid State Comm. **63**, 973 (1987).
- [15] V.J. Emery and S.A. Kivelson, Nature **374**, 434 (1995).
- [16] E.A. Kochetov and V.S. Yarunin, Phys. Rev. B **56**, 2703 (1997).
- [17] E.A. Kochetov and M. Mierzejewski, Phys. Rev. B **61**, 1580 (2000).